



IMAGEN-5.La submucosa se encontraba sustituida por tejido densamente colagenizado, pero era paucicelular; este tejido conectivo se introducía en el intersticio de la muscular propia, entre los haces musculares y se extendía perivesicalmente.

Immunohistochemistry (with differential diagnoses for comparison)							
	Inflammatory myofibroblastic tumour	postoperative spindle cell nodule	Sarcomatoid carcinoma	Stromal tumour of unknown malignant potential of the prostate	<u>Rhabdomyosarcoma</u>	<u>Leiomyosarcoma</u>	<u>Neurofibroma</u>
<u>ALK-1</u>	8/9 ¹ , 2/6 ⁵ , 3/4 ⁶ , 12/16 ⁷ , 12/26 ¹⁰		0/5 ¹ , 0/8 ⁷	0/2 ⁷	0/2 ¹	0/2 ¹ , 0/5 ⁷	0/2 ¹
<u>MNF116, Cam5.2 or AE1/3</u>	8/9 ¹ , 2/10 ² , 5/15 ⁴ , 13/16 ⁷ , 31/33 ¹⁰	2/4 ⁴	5/5 ¹ , 2/3 ⁴ , 6/8 ⁷	0/2 ⁷	0/2 ¹	1/2 ¹ , 3/8 ⁴ , 0/5 ⁷	0/2 ¹
<u>SMA</u>	8/9 ¹ , 3/8 ² , 5/8 ⁴ , 16/16 ⁷ , 23/24 ¹⁰	2/4 ⁴	4/5 ¹ , 2/3 ⁴ , 6/8 ⁷ , 6/8 ⁷	2/2 ⁷	0/2 ¹	2/2 ¹ , 3/7 ⁴ , 5/5 ⁷	0/2 ¹
<u>Desmin</u>	7/9 ¹ , 2/9 ² , 3/11 ⁴ , 12/15 ⁷ , 21/35 ¹⁰		2/5 ¹ , 0/3 ⁴ , 3/8 ⁷	2/2 ⁷	2/2 ¹	0/2 ¹ , 0/8 ⁴ , 3/5 ⁷	
<u>Calponin</u>	8/9 ¹					2/2 ¹	
<u>Caldesmon</u>	6/9 ¹					2/2 ¹	
<u>HHF35</u>	5/8 ⁴	2/3 ⁴	1/2 ⁴			1/5 ⁴	
<u>vimentin</u>	10/10 ² , 10/10 ⁴	4/4 ⁴	1/1 ⁴			7/7 ⁴	
<u>EMA</u>	2/8 ² , 4/8 ⁴	1/3 ⁴	1/2 ⁴			1/8 ⁴	
<u>p53</u>	7/10 ⁴ , 14/15 ⁷	3/4 ⁴	5/8 ⁷	0/2 ⁷		9/9 ⁴ , 3/5 ⁷	
<u>S-100</u>	0/4 ⁴ , 2/15 ⁷	0/3 ⁴	0/8 ⁷	0/2 ⁷		0/2 ⁴ , 0/5 ⁷	
<u>CD34</u>	0/15 ⁷		0/8 ⁷	0/2 ⁷		1/5 ⁷	
<u>CD117</u>	1/16 ⁷		1/8 ⁷	0/2 ⁷		1/5 ⁷	

TABLA-I. Tomado de: David N Louis, Herbert W Virgin IV, and Sylvia L Asa (2011) "Next-Generation" Pathology and Laboratory Medicine. Archives of Pathology & Laboratory Medicine: December 2011, Vol. 135, No. 12, pp. 1531-1532.